

<http://zoobank.org/urn:lsid:zoobank.org:pub:61513C30-CCAB-4EAA-A0FB-F61ED2BCA008>
DOI: 10.24412/2226-0773-2021-10-1-23-55

**New data on the distribution and biology of the longhorn beetles
from Taiwan (Coleoptera: Cerambycidae)**

W. Grosser¹, R. Ambrus²

¹U Střelnice 17, CZ-746 01 Opava, Czech Republic
e-mail: walterg@atlas.cz

²Tmukovo nám. 1112/1, CZ-152 00 Prague 5, Czech Republic
e-mail: ambrus@centrum.cz

Key words: Coleoptera, Cerambycidae, faunistics, biology, zoogeography, Taiwan.
Abstract: New data on the distribution and biology of the longhorn beetles occurring in Taiwan are presented together with a list of 156 species and subspecies representing 92 genera and 6 subfamilies that were collected during two entomological expeditions conducted in May 2012 and May/June 2013. New localities of 69 taxa recorded so far only from Taiwan are given. *Euryclytosemia nomurai* (Gressitt, 1951) and *Sybra botelensis* Breuning et K. Ohbayashi, 1966 are recorded from Lutao Isl. for the first time.

Introduction

Taiwan is a mountainous island in the western Pacific. The Tropic of Cancer runs through the middle part of Taiwan. The highest peak in the Backbone Mountains (or the Central Ridge) of this island is 3952 m. Two monsoon systems influence the climate in Taiwan, the summer monsoon from the south-western direction and the winter monsoon from the north-eastern direction. The warm and humid environment, long altitudinal gradients, different monsoon systems and complex topography cause the highly diverse flora and vegetation in Taiwan. According to the Flora of Taiwan (Huang at al., 1994-2003), there are 235 families, 1419 genera and 4340 species within an area of about 36 000 km². Among the 4340 species, 262 are naturalized and 1069 are endemic. Forests are the most dominant vegetation types in Taiwan, covering almost 60% of the land (www.forest.gov.tw). More than half of the natural forests are evergreen broad-leaved

forests and there are 463 evergreen broad-leaved woody species in the flora of Taiwan.

A great variety of habitats and natural vegetation is reflected in a great number of associated insect, many of them endemic to the island. The longhorn beetle fauna of Taiwan has been studied intensively over the past several decades, but additional taxa still await discovery. A total of approximately 806 Cerambycidae species and subspecies are now known from Taiwan out of which 508 are recorded so far only from this region.

Material and methods

A survey of the Cerambycidae of Taiwan was conducted out during years 2012 and 2013. The material for this study was collected by the first author and other collectors from different localities in Taiwan and is deposited in the personal collections.

The most effective standard methods for collecting beetles, such as attracting them to artificial light sources, shaking them into an entomological umbrella, sweep petting and analyses of the inhabited material, were used during the field research. The adults were determined by Carolus Holzschuh (Villach, Austria), Petr Viktora (Kutná Hora, Czech Republic), Tomáš Tichý (Opava, Czech Republic) and by the authors.

The following abbreviations of collectors are used in the text:

WG - Walter Grosser, Opava, Czech Republic

PK - Pavel Kučera, Liberec, Czech Republic

FČ - František Černý, Liberec, Czech Republic

Results

Family VESPERIDAE Subfamily Philinae

Tribe Philini

1) *Philus antennatus* (Gyllenhal, 1817)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 1 adult (WG).

Distribution: Known from SE Asia, China (Anhui, Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Hebei, Henan, Hongkong, Hubei, Hunan, Kiangsu, Kiangsi, Shaanxi, Shantung, Chekiang) and Taiwan.

2) *Philus pallescens ssp. pallescens* Bates, 1866

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 1 adult (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Pingtung prov., Kenting env., 21°56'52.21"N 120°46'57.23"E, 36 m, 4.6.2013, 1 adult (WG).

Distribution: Known from Laos, continental China (Fukien, Kwantung, Kweichow, Kwangsi, Henan, Hongkong, Hunan, Kiangsi, Sichuan, Shaanxi, Chekiang) and Taiwan.

Family DISTENIIDAE Subfamily Disteniinae

Tribe Cyrtonopini

3) *Cyrtonops asahinai* Mitono, 1947

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

W. Grosser, R. Ambrus

Family CERAMBYCIDAE
Subfamily Prioninae

Tribe Aegosomatini

4) *Aegolipton sauteri* (Lameere, 1913)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 12 adults (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 4 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 4 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

5) *Aegosoma sinicum ssp. sinicum* White, 1853

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 1 adult (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 2 adults (WG).

Distribution: Known from Russia (Far East), North Korea, South Korea, SE Asia, China (Anhui, Peking, Fukien, Gansu, Kwantung, Kweichow, Hainan, Hebei, Heilongjiang, Henan, Hubei, Hunan, Kiangsu, Kiangsi, Kirin, Kiangsi, Liaoning, Nei Mongol, Sichuan, Shanghai, Shandong, Shanxi, Tianjin, Tibet, Yunnan, Chekiang) and Taiwan.

Tribe Anacolini

6) *Sarmyds antennatus* Pascoe, 1867

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG).

Distribution: Known from India (Uttarakhand, Uttar Pradesh, Sikkim, Darjeeling), Bhutan, Nepal, SE Asia, China (Fukien, Kwantung, Kwangsi, Hainan, Hunan, Kiangsu, Shaanxi, Yunnan) and Taiwan.

Tribe Eurypodini

7) *Eurypoda antennata* Saunders, 1853

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 2 adults (WG).

Distribution: Known from China (Anhui, Kwantung, Kweichow, Kwangsi, Henan, Hongkong, Hainan, Hubei, Hunan, Kiangsu, Kiangsi, Sichuan, Shanghai, Tianjin, Tsianghai, Yunnan, Chekiang) and Taiwan.

W. Grosser, R. Ambrus

Tribe Macrotomini

8) *Bandar pascoei* ssp. *formosae* (Gressitt, 1938)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 3 adults (WG, PK, FČ).

Distribution: Known from Japan (Ryukyu Isl.) and Taiwan.

Tribe Prionini

9) *Prionus scabripunctatus* Hayashi, 1971

Material: Taiwan, Taichung prov., 15 km W Lishan, 24.25830°N 121.21086°E, 1413 m, 25.5.2012, 1 adult (PK).

Distribution: This taxon is so far known only from Taiwan.

10) *Priotyrranus closteroides* ssp. *closteroides* (Thomson, 1877)

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG).

Distribution: Known from Vietnam, China (Anhui, Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Henan, Hongkong, Hubei, Hunan, Kiangsu, Kiangsi, Sichuan, Liaoning, Nei Mongol, Sichuan, Shaanxi, Tianjin, Yunnan, Chekiang) and Taiwan.

Subfamily Lepturinae

Tribe Lepturini

11) *Anoploderomorpha formosana* (Matsushita, 1933)

Material: Taiwan, Nantou prov., 15 km of Wushe, Chingching Farm, 24.06680°N 121.16770°E, 2000 m, 13.6.2013, 1 adult (WG).

Distribution: Known from China (Hainan) and Taiwan.

12) *Anoploderomorpha izumii* (Tamanuki et Mitono, 1939)

Material: Taiwan, Taitung prov., Siangyang, 23.24774°N 120.98602°E, 2342 m, 31.5.-1.6.2013, 1 adult (FČ).

Distribution: Known from continental China (Fukien, Kwantung, Yunnan, Chekiang) and Taiwan.

13) *Leptura auratopilosa* (Matsushita, 1931)

Material: Taiwan, Taichung prov., 15 km W Lishan, 24.25830°N 121.21086°E, 1413 m, 25.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

14) *Nanostrangalia semichujoi* Hayashi, 1974

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 2 adults (WG).

W. Grosser, R. Ambrus

Distribution: This taxon is so far known only from Taiwan.

15) *Parastrangalis phantoma* Holzschuh, 1999

Material: Taiwan, Taichung prov., 15 km W Lishan, 24.25830°N 121.21086°E, 1413 m, 25.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

16) *Parastrangalis lateristriata* (Tamanuki et Mitono, 1939)

Material: Taiwan, Taitung prov., 20 km NW Litao, 23.24724°N 120.99052°E, 2120 m, 22.5.2012, 2 adults (WG); Taiwan, Taitung prov., Siangyang, 23.24774°N 120.98602°E, 2342 m, 31.5.-1.6.2013, 1 adult (WG).

Distribution: Known from SE Asia and Taiwan.

17) *Parastrangalis subapicalis* (Gressitt, 1935)

Material: Taiwan, Taitung prov., 20 km NW Litao, 23.24724°N 120.99052°E, 2120 m, 22.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

Tribe Rhagiini

18) *Pidonia (Mumon) sacrosancta* Kuboki, 1995

Material: Taiwan, Taitung prov., 20 km NW Litao, 23.24724°N 120.99052°E, 2120 m, 22.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

Subfamily Cerambycinae

Tribe Asemini

19) *Cephalallus oberthueri* Sharp, 1905

Material: Taiwan, Yilan prov., Tuleng env., 24.34637°N 121.31773°E, 1885 m, 10.5.2012, 1 adult (WG).

Distribution: Known from India (Arunachal Pradesh), continental China (Fukien, Kwangsi, Hubei, Kiangsi, Tibet, Yunnan) and Taiwan.

20) *Cephalallus unicolor ssp. unicolor* (Gahan, 1906)

Material: Taiwan, Taitung prov., Siangyang, 23.24774°N 120.98602°E, 2342 m, 31.5.-1.6.2013, 1 adult (WG).

Distribution: Known from North Korea, South Korea, Japan, SE Asia, China (Fukien, Kwantung, Kweichow, Hainan, Henan, Hongkong, Hubei, Hunan, Kiangsu, Kirin, Kiangsi, Sichuan, Yunnan, Chekiang) and Taiwan.

W. Grosser, R. Ambrus

Tribe Callichromatini

21) *Aphrodisium faldermannii* ssp. *horishaense* (Kano, 1933)

Material: Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

23) *Aphrodisium sauteri* (Matsushita, 1933)

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG).

Distribution: Known from continental China (Anhui, Fukien, Kwangsi, Henan, Hunan, Kiangsi, Shandong, Sichuan, Chekiang) and Taiwan.

24) *Embrikstrandia unifasciata* (Ritsema, 1897)

Material: Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG).

Distribution: Known from SE Asia, China (Anhui, Fukien, Gansu, Kwantung, Kwangsi, Hainan, Henan, Hongkong, Hubei, Hunan, Kiangsi, Sichuan, Shanxi, Chekiang) and Taiwan.

Tribe Callidiopini

25) *Ceresium elongatum* ssp. *elongatum* Matsushita, 1933

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 2 adults (WG); Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG).

Distribution: Known from Japan (Ryukyu Isl.), China (Hongkong) and Taiwan.

26) *Ceresium flavipes* (Fabricius, 1792)

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 6 adults (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 5 adults (WG).

Distribution: Known from Afrotropical Region, Australian Region, Neotropical Region, India, Pakistan, SE Asia, continental China (Kwantung, Hongkong) and Taiwan.

27) *Ceresium longicorne* Pic, 1926

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 2 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 8.6.2013, 1 adult (FČ).

Distribution: Known from South Korea, Japan, SE Asia, continental China (Hubei, Hongkong, Kiangsi) and Taiwan

28) *Ceresium sinicum* ssp. *sinicum* White, 1855

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 5 adults (WG).

Distribution: Known from South Korea, Japan, SE Asia, China (Peking, Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Henan, Hubei, Hunan, Kiangsu, Kiangsi, Sichuan, Shaanxi, Shanghai, Tibet, Yunnan, Chekiang) and Taiwan.

29) *Ceresium subuniforme* Schwarzer, 1925

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 3 adults (WG); Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 1 adult (FČ).

Distribution: This taxon is so far known only from Taiwan.

30) *Parasalpina kojimai* Hayashi, 1962

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 1 adult (WG).

Distribution: Known from Japan (Ryukyu: Iriomote Isl.) and Taiwan.

31) *Stenodryas clavigera* ssp. *impuncticollis* Hayashi, 1974

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 1 adult (WG); Taiwan, Taipei prov., Pinglin env., 24.97280°N 121.77214°E, 436 m, 16.-19.6.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

Tribe Cerambycini

32) *Gibbocerambyx maculicollis* (Matsushita, 1933)

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 3 adults (WG); Taiwan, Nantou prov., 5 km NE Shuili, 23.71029°N 120.87933°E, 900 m, 23.5.2012, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 1 adult (FC).

Distribution: This taxon is so far known only from Taiwan.

33) *Hemadius oenochrous* Fairmaire, 1889

Material: Taiwan, Yilan prov., Ssuchi (Siji) env., 24.50907°N 121.43788°E, 675 m, 26.5.2012, 3 adults on trunk of *Prunus serrulata* (WG, PK).

Distribution: Known from SE Asia, continental China (Anhui, Fukien, Kwangsi, Hubei, Hunan, Kiangsi, Sichuan, Tibet, Yunnan, Chekiang) and Taiwan.

34) *Lachnopterus socius* Gahan, 1891

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 16.5.2012, 1 adult (WG).

Distribution: Known from SE Asia and Taiwan.

35) *Margites fulvidus* (Pascoe, 1858)

Material: Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 2 adults (WG).

Distribution: Known from North Korea, South Korea, Japan, continental China (Fukien, Kwantung, Kweichow, Henan, Hubei, Hunan, Kiangsi, Sichuan, Shaanxi, Yunnan) and Taiwan.

36) *Massicus trilineatus* (Pic, 1933)

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 2 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 3 adults (WG, FC).

Distribution: Known from India (Arunachal Pradesh), SE Asia, China (Fukien, Kweichow, Kwangsi, Hainan, Kiangsi, Yunnan) and Taiwan.

37) *Nadezhdiella cantori* (Hope, 1845)

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (PK).

Distribution: Known from Thailand, China (Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Henan, Hongkong, Hubei, Hunan, Kiangsu, Kiangsi, Sichuan, Shaanxi, Yunnan, Chekiang) and Taiwan.

38) *Neoplocaederus bicolor* (Gressitt, 1942)

Material: Taiwan, Hsinchu prov., 20 km NW Neiwan, 24.69140°N 121.31076°E, 1563 m, 28.5.2013, 1 adult (PK).

Distribution: Known from China (Kweichow, Hainan, Hebei, Hubei, Hunan, Kiangsi, Shaanxi, Tibet, Yunnan) and Taiwan.

39) *Pseudopachydissus taiwanensis* Hayashi, 1992

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 2 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

40) *Trachylophus sinensis* Gahan, 1888

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG).

Distribution: Known from Vietnam, Myanmar, China (Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Hongkong, Hubei, Hunan, Kiangsi, Sichuan, Chekiang) and Taiwan.

41) *Trirachys indutus* (Newman, 1842)

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 16.5.2012, 1 adult (WG).

Distribution: Known from SE Asia, China (Anhui, Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Hongkong, Kiangsi, Chekiang) and Taiwan.

42) *Xoanodera maculata* Schwarzer, 1925

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 1 adult (WG); Taiwan, Kaoshiung prov., Baolai env., 21.10487°N 120.72800°E, 845 m, 10.6.2013, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (FČ).

W. Grosser, R. Ambrus

Distribution: Known from China (Fukien, Kwangsi, Hainan, Hunan, Sichuan, Yunnan) and Taiwan.

Tribe Cleomenini

43) *Cleomenes auricollis* Kano 1933

Material: Taiwan, Taitung prov., 20 km NW Litao, 23.24724°N 120.99052°E, 2120 m, 22.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

44) *Dere kirai* Mitono, 1943

Material: Taiwan, Hsinchu prov., 20 km NW Neiwan, 24.69140°N 121.31076°E, 1563 m, 15.6.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

Tribe Clytini

45) *Chlorophorus annularis* (Fabricius, 1787)

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 8.6.2013, 1 adult (FČ).

Distribution: Known from South Africa, Brazil, USA, Australia, Pakistan, India, Nepal, Japan, South Korea, SE Asia, China (Anhui, Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Hebei, Henan, Hongkong, Hubei, Hunan, Kiangsu, Kirin, Kiangsi, Liaoning, Sichuan, Shaanxi, Tibet, Yunnan, Chekiang) and Taiwan.

46) *Chlorophorus annulatus* (Hope, 1831)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 15 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 2 adults (WG); Taiwan, Taitung prov., 20 km NW Litao, 23.24724°N 120.99052°E, 2120 m, 22.5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 42 adults (WG); Taiwan, Taipei prov., Pinglin env., 24.97280°N 121.77214°E, 436 m, 16.-19.6.2013, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 1 adult (WG).

Distribution: Known from Japan, SE Asia, continental China (Fukien, Kwantung, Hubei, Kiangsi, Shaanxi, Chekiang) and Taiwan.

47) *Chlorophorus anticemaculatus* Schwarzer, 1925

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 2 adults (WG, FČ).

Distribution: This taxon is so far known only from Taiwan.

48) *Chlorophorus chiuae* Nakamura, 1974

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 3 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

49) *Chlorophorus kanekoi* Matsushita, 1941

Material: Taiwan, Nantou prov., Heshhe env., 23°33'44N 120°52'03E, 917 m, 23.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

50) *Chlorophorus cf. oppositus* (Chevrolat, 1863)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 3 adults (WG).

Distribution: Known from South continental China, Laos, Vietnam and Taiwan.

51) *Chlorophorus probsti* Holzschuh, 1989

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 2 adults (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

52) *Chlorophorus verus* Holzschuh, 1998

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 11 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

53) *Chlorophorus viridulus* Kano, 1933

Material: Taiwan, Chiayi prov., Shitzulu (Shizhuo), 23°28'N 120°41'E, 1350 m, 23.5.2012, 2 adults (WG); Taiwan, Nantou prov., Heshhe env., 23°33'44N 120°52'03E, 917 m, 23.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

54) *Chlorophorus taihorensis* Schwarzer, 1925

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 8 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

55) *Demonax bidenticornis* Matsushita, 1974

Material: Taiwan, Nantou prov., Heshe env., 23°33'44N 120°52'03E, 917 m, 23.5.2012, 2 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

56) *Demonax bimaculicollis* (Schwarzer, 1925)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG).

Distribution: Known from China (Hainan) and Taiwan.

57) *Demonax okunii* Mitono, 1942

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 3 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

58) *Demonax substitutus* Gressitt, 1934

Material: Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

59) *Perissus kankauensis* Schwarzer, 1925

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 18 adults (WG, FČ).

Distribution: This taxon is so far known only from Taiwan.

60) *Xylotrechus atronotatus* ssp. *atronotatus* Pic, 1917

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 22 adults (WG, FČ).

Distribution: Known from China (Fukie, Hainan, Kirin, Liaoning) and Taiwan.

61) *Xylotrechus grayii* ssp. *grayii* (White, 1855)

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 1 adult (WG).

Distribution: Known from South Korea, Japan (Mariana Isl.), continental China (Fukien, Gansu, Kwantung, Kweichow, Hebei, Henan, Hubei, Hunan, Kiangsu, Sichuan, Shaanxi, Northeast Territory, Shandong, Tibet, Yunnan) and Taiwan.

62) *Xylotrechus magicollis* (Fairmaire, 1888)

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 18 adults (WG, FČ).

Distribution: Known from India (Assam), Myanmar, Laos, Thailand, China (Fukien, Kwangsi, Hainan, Hebei, Hubei, Hunan,

W. Grosser, R. Ambrus

Kiangsi, Sichuan, Yunnan, Chekiang) and Taiwan.

Tribe Oabriini

63) *Pseudiphra apicalis* (Schwarzer, 1925)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG).

Distribution: Known from Japan, SE Asia and Taiwan.

64) *Uenobrium piceorubrum* (Hayashi, 1971)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

Tribe Oemini

65) *Nortia carinicornis* ssp. *carinicornis* Schwarzer, 1925

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 2 adults (WG).

Distribution: Known from Japan (Ryukyu Isl.), continental China (Fukien, Kwangsi) and Taiwan.

66) *Oplatocera (Epioplatocera) mitonoi* Hayashi, 1981

Material: Taiwan, Taipei prov., Pinglin env., 24.97280°N 121.77214°E, 436 m, 16.-19.6.2013, 1 adult (WG).

Distribution: Known from continental China (Chekiang) and Taiwan.

Tribe Pseudolepturini

67) *Erythrurus formosanus* Bates, 1866

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

Tribe Purpuricenini

68) *Dicelosternus corallinus* Gahan, 1900

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 11 adults (WG, FČ).

Distribution: Known from China (Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Hubei, Hunan, Kiangsi, Chekiang) and Taiwan.

Tribe Stenhomalini

69) *Stenhomalus fenestratus* White, 1855

Material: Taiwan, Nantou prov., 5 km NE Shuili, 23.71029°N 120.87933°E, 900 m, 23.5.2012, 1 adult (WG).

Distribution: Known from India, Nepal, SE Asia, continental China (Fukien, Kwantung, Hongkong, Northeast Territory, Northern Territory, Shaanxi, Sichuan) and Taiwan.

Tribe Stenopterini

70) *Merionoeda (Ocytasia) formosana ssp. formosana* Heller, 1924

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 2 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 14 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

71) *Microdebilissa testacea* Matsushita, 1933

Material: Taiwan, Nantou prov., Heshe env., 23°33'44N 120°52'03E, 917 m, 23.5.2012, 11 adults (WG); Taiwan, Yilan prov., Ssuchi (Siji) env., 24.50907°N 121.43788°E, 675 m, 26.5.2012, 1 adult (WG).

Distribution: Known from Nepal, continental China (Yunnan) and Taiwan.

Tribe Thraniini

72) *Thranium multinotatus ssp. signatus* Schwarzer, 1925

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (PK); Taiwan, Yilan prov., Chingshui env., 29.-30.5.2013, 24.62943°N 121.57557°E, 257 m, 2 adults (WG, FČ).

Distribution: Known from SE Asia, China (Fukien, Hainan, Chekiang) and Taiwan.

73) *Thranium rufescens ssp. formosanus* Schwarzer, 1925

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 2 adults (WG, PK).

Distribution: This taxon is so far known only from Taiwan.

W. Grosser, R. Ambrus

Tribe Tillomorphini

74) *Epipedocera rollei* Pic, 1910

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6. 2013, 69 adults (WG, FČ); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 8.6.2013, 1 adult (FČ).

Distribution: This taxon is so far known only from Taiwan.

Tribe Xystrocerini

75) *Leptoxenus bimaculatus* (Matsushita, 1933)

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

76) *Xystrocera globosa* (Olivier, 1795)

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG).

Distribution: Known from Afrotropical Region, Australian Region, Naotropical Region, Egypt, Israel, Pakistan, India, Nepal, Bhutan, North Korea, South Korea, Japan, SE Asia, China (Anhui, Fukien, Gansu, Kwantung, Kweichow, Kwangsi, Hainan, Hebei, Henan, Hubei, Hunan, Kiangsu, Kiangsi, Sichuan, Shaanxi, Shanghai, Shandong, Yunnan, Chekiang) and Taiwan.

Subfamily Lamiinae

Tribe Acanthocinini

77) *Euryclytosemia nomurai* (Gressitt, 1951)

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 1 adult (PK); Taiwan, Lutaο Isl., 6.-7.6.2013, 1 adult (WG).

Distribution: Known from Japan (Ryukyu Isl.) and Taiwan (Lanyu and Lutaο Isl.).

Remark: The species is recorded from Lutaο Isl. for the first time.

78) *Rondibilis horiensis* ssp. *horiensis* Kano, 1933

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6. 2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

79) *Rondibilis semielongata* Hayashi, 1974

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 45 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 2 adults (WG); Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 2 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

80) *Rondibilis taiwana* Hayashi, 1974

Material: Taiwan, Taitung prov., Siangyang, 23.24774°N 120.98602°E, 2342 m, 31.5.-1.6.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

Tribe Agapanthiini

81) *Pothyne albolineata* Matsushita, 1933

Material: Taiwan, Nantou prov., 5 km NE Shuili, 23.71029°N 120.87933°E, 900 m, 23.5. 2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

82) *Pothyne formosana ssp. formosana* Schwarzer, 1925

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG).

Distribution: Known from continental China (Kwantung) and Taiwan.

83) *Pothyne formosana ssp. nanshanchina* Takakuwa et Kusama, 1979

Material: Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 2 adults (WG); Taiwan, Nantou prov., Sun Moon Lake env. 23.82303°N 120.94062°E, 800 m, 12.6.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

84) *Pothyne variegata ssp. variegata* Thomson, 1864

Material: Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 2 adults (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 1 adult (WG).

Distribution: Known from N India (Sikkim, Darjeeling), Nepal, SE Asia, continental China (Hunan, Hongkong, Yunnan) and Taiwan.

W. Grosser, R. Ambrus

85) *Pseudocalamobius pubescens* Hasegawa, 1987

Material: Taiwan, Nantou prov., 15 km of Wushe, Chingching Farm, 24.06680°N 121.16770°E, 2000 m, 13.6.2013, 1 adult (WG); Taiwan, Taitung prov., Siangyang, 23.24774°N 120.98602°E, 2342 m, 31.5.-1.6.2013, 1 adult (FC).

Distribution: This taxon is so far known only from Taiwan.

Tribe Ancylonotini

86) *Palimna palimnoides* (Schwarzer, 1925)

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6. 2013, 2 adults (PK); Taiwan, Taitung prov., Siangyang, 23.24774°N 120.98602°E, 2342 m, 3.6.2013, 1 adult (FC).

Distribution: Known from India (Sikkim, Darjeeling), Nepal, Bhutan, Laos, China (Fukien, Kwangsi, Hainan, Yunnan) and Taiwan.

Tribe Apodasyini

87) *Cylindilla formosana* (Gressitt, 1951)

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

88) *Euseboides matsudai ssp. matsudai* Gressitt, 1938

Material: Taiwan, Pingtung prov., Kenting env., 21°56'52.21"N 120°46'57.23"E, 36 m, 19.5.2012, 1 adult (WG).

Distribution: Known from Japan and Taiwan.

89) *Falsoterinaea rufipennis* (Matsushita, 1933)

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 6 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

90) *Mimectatina murakamii* Hayashi, 1978

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan (Lanyu Isl.).

91) *Penthides flavus ssp. flavus* Matsushita, 1933

Material: Taiwan, Nantou prov., Sun Moon Lake env.,

W. Grosser, R. Ambrus

23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

92) *Rhodopina formosana* (Breuning, 1954)

Material: Taiwan, Yilan prov., 20 km S Yilan, Yulan, 24.70017°N 121.64717°E, 95 m, 8.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

Tribe Apomecynini

93) *Apomecyna histrio* ssp. *histrio* (Fabricius, 1792)

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 4 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 1 adult (WG).

Distribution: Known from Russia (E Siberia), Pakistan, Nepal, India (Arunachal and Uttar Pradesh, Sikkim, Darjeeling), Mongolia, S Korea, Japan (Ryukyu Isl.), SE Asia, China (Hainan, Yunnan, Kweichow) and Taiwan.

94) *Doliops similis* Miwa et Mitono, 1933

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 1 adult (PK).

Distribution: This taxon is so far known only from Taiwan (Lanyu Isl.).

95) *Ropica dorsalis* Schwarzer, 1925

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG); Taiwan, Nantou prov., Sun Moon Lake env. 23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG).

Distribution: Known from India (Sikkim, Darjeeling, Uttar Pradesh), Nepal, Bhutan, Japan, SE Asia, China (Kwantung, Kwangsi, Hainan, Kiangsu) and Taiwan.

96) *Ropica formosana* Bates, 1866

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 3 adults (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N

W. Grosser, R. Ambrus

121.15394°E, 403 m, 31.5. 2013, 2 adults (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 2 adults (WG); Taiwan, Kaoshiung prov., Baolai env., 21.10487°N 120.72800°E, 845 m, 10.6.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

97) *Ropica fuscolaterimaculata* Hayashi, 1974

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 4 adults (WG).

Distribution: This taxon is so far known only from Taiwan (Lanyu Isl.).

98) *Ropica honesta* Pascoe, 1865

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 2 adults (WG).

Distribution: Known from Nepal, Japan, SE Asia, China (Hongkong) and Taiwan.

99) *Sybra baculina* Bates, 1866

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG).

Distribution: Known from Japan (Ryukyu Isl.), China (Hongkong) and Taiwan.

100) *Sybra bioculata* ssp. *quadrinotata* Schwarzer, 1925

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., N 23.08212° E 121.28433°, 248 m, 2.6.2013, 1 adult (WG).

Distribution: Known from Vietnam, continental China (Kiangsu, Yunnan, Chekiang) and Taiwan.

101) *Sybra botelensis* Breuning et K. Ohbayashi, 1966

Material: Taiwan, Lanyu Isl. 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 2 adults (WG). Taiwan, Lutao Isl., 6.-7.6.2013, 9 adults (WG).

Distribution: This taxon is so far known only from Taiwan (Lanyu and Lutao Isl.).

Remark: The species is recorded from Lutao Isl. for the first time.

102) *Sybra flavostriata* ssp. *flavostriata* Hayashi, 1968

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m,

W. Grosser, R. Ambrus

18.5.2012, 1 adult (WG).

Distribution: Known from Japan (Ryukyu Isl.) and Taiwan.

103) *Sybra ordinata ssp. ordinata* Bates, 1873

Material: Taiwan, Yilan prov., Chingshui env., 24.62943°N 121.57557°E,

257 m, 29.-30.5.2013, 1 adult (WG); Taiwan, Yilan prov., Chingshui env., 24.62943°N 121.57557°E, 257 m, 29.-30.5.2013, 3 adults (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG).

Distribution: Known from China (Fukien, Kwangsi, Hainan, Kiangsu, Kiangsi, Shanghai) and Taiwan.

104) *Sybra pascoei ssp. taiwanella* Gressitt, 1951

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 2 adults (WG). Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 3 adults (PK).

Distribution: Known from Japan (Ryukyu Isl.) and Taiwan.

105) *Sybra posticalis* (Pascoe, 1858)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 1 adult (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 3 adults (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 4 adults (WG).

Distribution: Known from China (Hainan, Hongkong) and Taiwan.

Tribe Batocerini

106) *Apriona rugicollis ssp. rugicollis* Chevrolat, 1852

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 5 adults (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 2 adults (WG, FČ); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 3 adults (WG, FČ); Taiwan, Kaoshiung prov., Baolai env.,

W. Grosser, R. Ambrus

21.10487°N 120.72800°E, 845 m, 10.6.2013, 3 adults (WG, FČ); Taiwan, Taitung prov., Siangyang, 23.24774°N 120.98602°E, 2342 m, 31.5.-1.6.2013, 1 adult (FČ); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 8.6.2013, 2 adults (FČ).

Distribution: Known from Russia (Far East), North Korea, South Korea, China (Anhui, Fukien, Gansu, Kwantung, Kweichow, Kwangsi, Hainan, Hebei, Henan, Hongkong, Hubei, Hunan, Kiangsu, Kiangsi, Liaoning, Sichuan, Shaanxi, Shandong, Shanxi, Tibet, Yunnan, Chekiang) and Taiwan.

107) *Batocera lineolata* Chevrolat, 1852

Material: Taiwan, Hsinchu prov., 20 km NW Neiwan, 24.69140°N 121.31076°E, 1563 m, 28.5.2013, 2 adults (WG, PK); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG).

Distribution: Known from India (Sikkim, Darjeeling), Japan, South Korea, China (Anhui, Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Hebei, Hubei, Kiangsu, Kiangsi, Sichuan, Shaanxi, Yunnan, Chekiang) and Taiwan.

Tribe Dorcaschematini

108) *Microlenecamptus biocellatus* (Schwarzer, 1925)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 1 adult (WG).

Distribution: Known from continental China (Anhui, Kwantung, Hunan) and Taiwan.

109) *Olenecamptus bilobus* ssp. *bilobus* (Fabricius, 1801)

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 4 adults (WG); Taiwan, Lutao Isl., 6.-7.6.2013, 1 adult (WG).

Distribution: Known from Afrotropical Region, Australian Region, Pakistan, Nepal, India (Sikkim, Darjeeling, Uttar, Himachal and Arunachal Pradesh), SE Asia, China (Fukien, Kwantung, Kwangsi, Hainan, Hebei, Hongkong, Liaoning, Sichuan, Yunnan, Chekiang) and Taiwan (Lanyu and Lutao Isl.).

110) *Olenecamptus cretaceus* ssp. *marginatus* Schwarzer, 1925

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, ex larva, 5 adults ex larva from 18.7. to 5.8.2012 (WG); Taiwan, Taitung prov., 20 km N

W. Grosser, R. Ambrus

Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 18 adults ex larva from 12.7. to 15.9. 2013 (WG); Taiwan, Nantou prov., 5 km NE Shuili, 23.71029°N 120.87933°E, 900 m, 23.5.2012, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 26 adults ex larva from 12.7. to 15.9. 2013 (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 8.6.2013, 1 adult (FC).

Distribution: Known from SE Asia, continental China (Henan) and Taiwan.

111) *Olenecamptus formosanus* Pic, 1914

Material: Taiwan, Yilan prov., 20 km S Yilan, 24.70017°N 121.64717°E, 95 m, 8.5.2012, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 2 adults (WG).

Distribution: Known from S Korea, Japan, SE Asia, continental China (Hubei, Chekiang) and Taiwan.

112) *Olenecamptus taiwanus* Dillon et Dillon, 1948

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 5 adults (WG, FC).

Distribution: Known from Japan, China (Kwantung, Kwangsi, Hainan, Yunnan, Hongkong) and Taiwan.

Tribe Eunidiini

113) *Eunidia taiwanensis* Hayashi et Nara, 1992

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

Tribe Exocentrini

114) *Exocentrus brevisetosus* Gressitt, 1938

Material: Taiwan, Nantou prov., 5 km NE Shuili, 23.71029°N 120.87933°E, 900 m, 23.5.2012, 1 adult (WG).

Distribution: Known from SE Asia, continental China (Hubei, Yunnan) and Taiwan.

W. Grosser, R. Ambrus

115) *Exocentrus immaculatus* Gressitt, 1951

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

116) *Exocentrus pseudovaripeennis* Kusama et Tahira, 1925

Material: Taiwan, Nantou prov., Sun Moon Lake env. 23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG).

Distribution: Known from SE Asia and Taiwan.

117) *Exocentrus seriatomaculatus* Schwarzer, 1925

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 26 adults ex larva from 11.7. to 14.10.2012 (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 3 adults (WG).

Distribution: Known from SE Asia and Taiwan.

118) *Miaenia (Truncatomiaenia) botelensis* (Gressitt, 1951)

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 1 adult (PK); Taiwan, Lutao Isl., 6.-7.6.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan (Lanyu and Lutao Isl.).

Tribe Homonoeni

119) *Bumetopia lanshuana* Hayashi, 1974

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan (Lanyu Isl.).

120) *Bumetopia stolata* (Matsushita, 1931)

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 4 adults (WG).

Distribution: This taxon is so far known only from Taiwan (Lanyu Isl.).

Tribe Mesosini

121) *Agelasta (Dissosira) perplexa* Pascoe, 1858

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N

W. Grosser, R. Ambrus

120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 5 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 4 adults (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6. 2013, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 4 adults (WG).

Distribution: Known from Japan, South Korea (Gageodo Isl.), continental China (Fukien, Henan, Kiangsi, Liaoning, Chekiang) and Taiwan.

122) *Agelasta (Dissosira) tonkinea ssp. tonkinea* Pic, 1925

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult ex larva 28.1.2014 (WG).

Distribution: Known from Nepal, SE Asia and Taiwan.

123) *Coptops japonicus* Breuning, 1936

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult ex larva 5.9.2012 (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 2 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 1 adult (WG); Taiwan, Hsinchu prov., 20 km NW Neiwan, 24.69140°N 121.31076°E, 1563 m, 28.5.2013, 1 adult (FČ).

Distribution: This taxon is so far known only from Taiwan.

124) *Mesoereis bifasciata* (Pic, 1925)

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG); Taiwan, Hsinchu prov., 20 km NW Neiwan, 24.69140°N 121.31076°E, 1563 m, 28.5.2013, 1 adult (WG).

Distribution: Known from SE Asia, China (Fukien, Kwangsi, Hainan) and Taiwan.

125) *Mesosa (Aplocnemia) latifasciata* (White, 1858)

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG).

Distribution: Known from Vietnam, China (Fukien, Gansu, Kwantung, Kweichow, Kwangsi, Hainan, Hebei, Kiangsu, Kiangsi, Sichuan, Shanxi, Chekiang) and Taiwan.

Tribe Monochamini

126) *Acalolepta grossescapa* (Breuning 1942)

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

127) *Acalolepta (Dihammus) rusticator ssp. formosensis* Breuning et K. Ohbayashi, 1966

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 3 adults (WG).

Distribution: This taxon is so far known only from Taiwan (Lanyu and Lutaο Isl.).

128) *Acalolepta (Dihammus) rusticator ssp. rusticator* (Fabricius, 1801)

Material: Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 2 adults (WG).

Distribution: Known from SE Asia and Taiwan.

129) *Acalolepta sublusca ssp. maculihumera* (Matsushita, 1933)

Material: Taiwan, Hsinchu prov., 20 km NW Neiwan, 24.69140°N 121.31076°E, 1563 m, 15.6.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

130) *Anoplophora albopicta* (Matsushita, 1933)

Material: Taiwan, Taipei prov., Pinglin env., 24.97280°N 121.77214°E, 436 m, 16.-19.6.2013, 1 adult (PK).

Distribution: This taxon is so far known only from Taiwan.

131) *Anoplophora horsfieldii ssp. taiwanensis* (Hope, 1842)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6. 2013, 1 adult (PK).

Distribution: This taxon is so far known only from Taiwan.

132) *Anoplophora macularia* (Thomson, 1865)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 7 adults (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Nantou prov., Sun Moon Lake env. 23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG); Taiwan, Yilan prov., Ssuchi (Siji) env., 24.50907°N 121.43788°E, 675 m, 26. 5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 2 adults (WG, FC); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 8.6.2013, 1 adult (FC); Taiwan,

W. Grosser, R. Ambrus

Lutao Isl., 6.-7.6.2013, 2 adults (FČ).

Distribution: Known from Japan (Ryukyu Isl.), SE Asia, China (Fukien, Kwantung, Kwangsi, Hainan, Henan, Kiangsu, Sichuan, Chekiang, Northeast Territory) and Taiwan.

133) *Astynoscelis degener* (Bates, 1873)

Material: Taiwan, Taichung prov., 15 km W Lishan, 24.25830°N 121.21086°E, 1413 m, 25.5.2012, 1 adult (WG).

Distribution: Known from Russia (Far East), Japan, South Korea, continental China (Fukien, Gansu, Kwantung, Kweichow, Kwangs, Heilongjiang, Hubei, Hunan, Kiangsu, Kirin, Kiangsi, Nei Mongol, Sichuan, Shaanxi, Shandong, Yunnan, Chekiang) and Taiwan.

134) *Blepephaeus succinator* (Chevrolat, 1852)

Material: Taiwan, Nantou prov., 5 km NE Shuili, 23.71029°N 120.87933°E, 900 m, 23.5.2012, 1 adult (WG).

Distribution: Known from India (Arunachal Pradesh, Sikkim, Darjeeling), Nepal, SE Asia, China (Kwantung, Kwangsi, Hainan, Hongkong, Hunan, Kiangsu, Kiangsi, Sichuan, Shaanxi, Shanghai, Tibet, Yunnan, Chekiang) and Taiwan.

135) *Epepeotes ambigenus ssp. formosanus* Gressitt, 1951

Material: Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 1 adult (WG); Taiwan, Lutao Isl., 6.-7.6.2013, 2 adults (WG, FČ).

Distribution: This taxon is so far known only from Taiwan (Lanyu and Lutao Isl.).

136) *Pebblephaeus decoloratus* (Schwarzer, 1925)

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Nantou prov., 5 km NE Shuili, 23.71029°N 120.87933°E, 900 m, 23.5.2012, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 1 adult (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 2 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

137) *Psacotha hilaris ssp. hilaris* (Pascoe, 1857)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N

W. Grosser, R. Ambrus

121.28433°E, 248 m, 11.-12.5.2012, 2 adults (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 12 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 4 adults (WG, FČ); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 1 adult (WG); Taiwan, Kaoshiung prov., Baolai env., 21.10487°N 120.72800°E, 845 m, 10.6.2013, 3 adults (WG, FČ); Taiwan, Taitung prov., Siangyang, 23.24774°N 120.98602°E, 2342 m, 31.5.-1.6.2013, 1 adult (FČ).

Distribution: Known from France, Italy, Japan, South Korea, SE Asia, China (Anhui, Fukien, Gansu, Kwantung, Kweichow, Kwangsi, Hainan, Hebei, Henan, Hubei, Hunan, Kiangsu, Kiangsi, Northeast Territory, Sichuan, Shaanxi, Yunnan, Chekiang) and Taiwan.

138) *Uraecha angusta* (Pascoe, 1856)

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG).

Distribution: Known from SE Asia, continental China (Anhui, Pekung, Fukien, Kwantung, Kweichow, Kwangsi, Hebei, Henan, Hubei, Hunan, Kiangsu, Kiangsi, Sichuan, Shaanxi, Tibet, Chekiang) and Taiwan.

Tribe Phytoeciini

139) *Obereopsis kankauensis* (Schwarzer, 1925)

Material: Taiwan, Yilan prov., Ssuchi (Siji) env., 24.50907°N 121.43788°E, 675 m, 26.5.2012, 1 adult (WG).

Distribution: Known from continental China (Fukien, Sichuan) and Taiwan.

140) *Oberea flavipennis* Kurihara et N. Ohbayashi, 2007

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

141) *Oberea formosana* Pic, 1911

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 5 adults (WG); Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Kaoshiung prov., Baolai env., 21.10487°N 120.72800°E, 845 m, 10.6.2013, 1 adult (FČ).

Distribution: Known from Nepal, SE Asia, China (Anhui, Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Henan, Hubei, Hunan, Kiangsu, Kiangsi, Sichuan, Shaanxi, Chekiang) and Taiwan.

Tribe Pteropliini

142) *Desisa variabilis* (Schwarzer, 1925)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 16 adults (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 12 adults ex larva from 10.2. to 26.3.2014 (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 23 adults ex larva from 8.6 to 27.7.2013 (WG, FČ); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 2 adults (WG).

Distribution: This taxon is so far known only from Taiwan.

143) *Egesina (Nijjimaia) fuscolaterimaculata* Hayashi, 1971

Material: Taiwan, Chiayi prov., Shitzulu (Shizhuo), 23°28'N 120°41'E, 1350 m, 23.5.2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

144) *Pterolophia (Hylobrotus) annulata* (Chevrolat, 1845)

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 2 adults (WG); Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 3 adults (WG); Taiwan, Nantou prov., Sun Moon Lake env., 23.82303°N 120.94062°E, 800 m, 24.5.2012, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 2 adults (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 1 adult (WG).

Distribution: Known from Nepal, India (Sikkim, Darjeeling), S

W. Grosser, R. Ambrus

Korea, Japan, SE Asia, China (Fukien, Kwantung, Kweichow, Kwangsi, Hainan, Hebei, Henan, Hongkong, Hubei, Hunan, Kiangsu, Kiangsi, Sichuan, Shaanxi, Yunnan, Chekiang) and Taiwan.

145) *Pterolophia bigibbera* (Newman, 1842)

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Lanyu Isl., 22.06827°N 121.56625°E, 22 m, 15.-17.5.2012, 4 adults (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 2 adults (WG); Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 1 adult (WG); Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 2 adults (WG).

Distribution: Known from Japan (Ogasawara Isl.), Philippines and Taiwan.

146) *Pterolophia kaleea* ssp. *kaleea* (Bates, 1866)

Material: Taiwan, Taitung prov., 20 km N Taitung, 22.87112°N 121.16184°E, 424 m, 13.5.2012, 1 adult (WG); Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5.2012, 3 adults (WG).

Distribution: Known from Japan (Ryukyu Isl.), China (Kweichow, Hainan) and Taiwan.

147) *Pterolophia (Hylobrotus) latefascia* Schwarzer, 1925

Material: Taiwan, Yilan prov., Chingshui env., 24.62943°N 121.57557°E, 257 m, 29.-30.5.2013, 1 adult (WG).

Distribution: Known from Japan (Ryukyu Isl.) and Taiwan.

148) *Pterolophia laterialba* (Schwarzer, 1925)

Material: Taiwan, Taitung prov., 20 km N Taitung, N 22.87112° E 121.16184°, 424 m, 13.5.2012, 1 adult (PK).

Distribution: This taxon is so far known only from Taiwan.

149) *Pterolophia obscura* Schwarzer, 1925

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

150) *Pterolophia reduplicata* Gressitt, 1951

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 2.6.2013, 1 adult (WG).

Distribution: Known from continental China (Hubei) and Taiwan.

Tribe Saperdini

151) *Epiglenea comes* ssp. *formosana* (Schwarzer, 1925)

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

152) *Glenea chrysomaculata* Schwarzer, 1925

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG); Taiwan, Taitung prov., Wulu env., 23.12762°N 121.15394°E, 403 m, 31.5.2013, 1 adult (WG).

Distribution: Known from central China (Kweichow, Kwangsi) and Taiwan.

153) *Glenea lineata* ssp. *sauteri* Schwarzer, 1925

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 3 adults (WG).

Distribution: Known from continental China (Kwantung, Yunnan) and Taiwan.

154) *Glenea nanshanchiana* Hayashi, 1978

Material: Taiwan, Taitung prov., Taiyuan env., 23.08212°N 121.28433°E, 248 m, 11.-12.5. 2012, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

155) *Glenea relicta* ssp. *formosensis* Breuning, 1960

Material: Taiwan, Taitung prov., Schouchia env., 22.20477°N 120.86710°E, 259 m, 3.-5.6.2013, 1 adult (WG).

Distribution: This taxon is so far known only from Taiwan.

156) *Paraglenea swinhoei* ssp. *swinhoei* Bates, 1866

Material: Taiwan, Taitung prov., Schouchia env., 22.20670°N 120.86034°E, 315 m, 18.5.2012, 4 adults (WG); Taiwan, Chiayi prov., Shihtzulu (Shizhuo), 23°28'N 120°41'E, 1350 m, 23.5.2012, 1 adult (WG); Taiwan, Yilan prov., Chingshui env., 24.62943°N 121.57557°E, 257 m, 29.-30.5.2013, 4 adults (WG, FC).

Distribution: This taxon is so far known only from Taiwan.

Acknowledgements. We are very grateful to Carolus Holzschuh (Villach, Austria), Petr Viktora (Kutná Hora, Czech Republic) and Tomáš Tichý (Opava, Czech Republic) for determination and comments on the collected species.

REFERENCES

- Chang S.-C. 1960. New and unrecorded species of longicorn beetles from Taiwan (I). - Bulletin of the National Institut of Agricultural Science, Plant Pathology and Entomology 11: 1-7, pls 1, 2.
- Chang S.-C. 1978. New longicorn beetles of Taiwan (Coleoptera: Cerambycidae). - Journal of Agriculture and Forestry 27: 59-66.
- Chou W.-I. 2004. Taiwan Tianniu Tujian. Taipei, Maotouying Chubanshe: 408 pp. [Iconography of Longhorn Beetles in Taiwan. Taipei: Owl Press].
- Chou W.-I. 2008. The Atlas of Taiwanese Cerambycidae (second edition). Taipei, Owl Press, 408 pp.
- Chou W.-I., Ohbayashi N. 2007. Notes on the genus *Sinostrangalis* and *Metastrangalis* (Coleoptera, Cerambycidae). Studies on the Taiwanese Lepturinae. - Japanese Journal of Systematic Entomology. 13 (2): 225-240.
- Chou W.-I., Ohbayashi N. 2008. A new species of the genus *Caraphia* from Taiwan, with consideration of the status of the genus (Coleoptera, Cerambycidae). Studies on the Taiwanese Lepturinae, ii. - Entomological Review of Japan. 63: 135-144.
- Chou W.-I., Ohbayashi N. 2010. A New Genus and Four New Species of Taiwanese Lepturinae (Coleoptera, Cerambycidae). Studies on the Taiwanese Lepturinae, iii. - Japanese Journal of Systematic Entomology. 16 (2): 359-371.
- Chou W.-I., Chung Y.-T., Lin M.-Y. 2010. Description of *Eutetrappa lini* sp. nov. from Taiwan, China. (Coleoptera, Cerambycidae, Lamiinae, Saperdini). - Acta Zootaxonomica Sinica. 35 (2): 313-318.
- Chou W.-I., Ohbayashi N. 2014. Two New Species of Taiwanese Lepturini (Coleoptera: Cerambycidae). Studies on the Taiwanese Lepturinae, vi. - Japanese Journal of Systematic Entomology. 20 (2): 347-352.
- Chou W.-I., Wu W.-J. 2005. Studies on Cerambycidae of Taiwan and adjacent Regions. ii. Two new lepturine species of the genus *Pidonia* Mulsant (Coleoptera, Cerambycidae) from the subalpine zone of southern Taiwan. - Elytra. 33: 403-411.
- Danilevsky M.L. (editor) 2020. Catalogue of Palaearctic Coleoptera, Volume 6/1. Chrysomeloidea I (Vesperidae, Disteniidae, Cerambycidae). Updated and Revised Second Edition. Brill: 924 pages, 1 b/w illustration, 3 b/w maps.
- Drumont A., Ripaille C. 2019. Description of a new species belonging to the genus *Sarmyds* Pascoe from Taiwan island, China (Coleoptera, Cerambycidae, Prioninae) (6th contribution to the study of genus *Sarmyds* Pascoe, 1867). - Lambillionea. 119 (3): 158-166.
- Gressitt J.L. 1951. Longicorn beetles of China. Longicornia, Paris, 2: 1-667, 22 pls.
- Huang T.-C. and Editorial Committee of the Flora of Taiwan. (eds.). 1994-2003. Flora of Taiwan, Volume 1-6, 2nd edn. Editorial Committee of the Flora of Taiwan, Department of Botany, National Taiwan University, Taipei, Taiwan.
- Hasegawa M. 1987. Cerambycid beetles of the genus *Pseudocalamobius*

W. Grosser, R. Ambrus

- (Coleoptera) from Taiwan. - Kontyu. 55: 509-520.
- Heyrovský L. 1955b. Zwei neue Cerambyciden-Arten aus Taiwan (Coleoptera). - Annales Historico-Naturales Musei Hungarici Hungarici (n.s.) 6: 259-261.
- Holzschuh, C. 1989b: Beschreibung von 20 neuen Bockkäfern aus Thailand, Vietnam und Taiwan (Coleoptera, Cerambycidae). - Entomologica Basiliensia. 13: 361-390.
- Holzschuh C., Lin Y.-L. 2015. Beitrag zur Bockkaferfauna von Taiwan mit Beschreibung einer neuen Art, ii (Coleoptera, Cerambycidae). - Les Cahiers Magellanes. 20 (ns): 104-107.
- Kurihara T., Ohbayashi N. 2006. Two new species of the genus *Oberia* Dejean from Taiwan (Coleoptera, Cerambycidae). - Japanese Journal of Systematic Entomology. 12: 257-265.
- Makihara, H. 1978. New longicorn-beetles of the Ryukyus and Taiwan (Coleoptera: Cerambycidae). - Esakia. 12: 49-62.
- Nakamura Sh., Makihara H., Kurihara T., Yamasako J. 2014. Check-list of Longicorn-beetles of Taiwan. - Miscellaneous Report of the Hiwa Museum for the Natural History. 55: 1-278.
- Niisato T., Chou W.-I. 2018. Eight New Taxa and a New Record of the Genera *Obrium* Dejean and *Stenhomalus* A. White (Coleoptera: Cerambycidae) from China and Taiwan. - Japanese Journal of Systematic Entomology. 24 (1): 73-88.
- Shimomura T. 1979. New cerambycid beetles of the genus *Lemula* Bates from Taiwan. - Bulletin of the National Science Museum, Tokyo (Zoology). 5: 271-279.
- Shimomura T. 1993. A revision of the Taiwanese species of *Robustanoplodera* (Coleoptera, Cerambycidae, Lepturinae). - Elytra. 21: 227-234.
- Yamasako, J. 2017: A new species of the genus *Eupogoniopsis* (Coleoptera, Cerambycidae, Lamiinae) from Taiwan. - Special Bulletin of the Coleopterological Society of Japan. 1: 243-246.
- Yamasako J., Chou W.-I. 2014a. A new species of the genus *Callomecyna* from Taiwan (Coleoptera: Cerambycidae). Acta Entomologica Musei Nationalis Pragae 54 (1): 251-256.

Received: 13.12.2020

Accepted: 14.01.2021